SOLAR OBSERVATIONS

SOLAR AND SKY RADIATION MEASUREMENTS DURING MAY, 1927

By HERBERT H. KIMBALL, Solar Radiation Investigations

For a description of instruments and exposures and an account of the method of obtaining and reducing the measurements, the reader is referred to the Review for January, 1924, 52:42, January, 1925, 53:29, and July, 1925, 53:318.

From Table 1 it is seen that solar-radiation intensities averaged above the May normals at all three stations. At Washington a noon intensity of 1.45 gram-calories per minute per square centimeter obtained on the 27th equals the highest noon intensity heretofore obtained at that station in May.

Table 2 shows a deficiency in the total solar radiation received on a horizontal surface from the sun and sky at all three stations for which normals have been determined. The deficiency is pronounced at Washington and Madison and is due to unusually cloudy conditious.

Skylight-polarization measurements were mad at Washington on only one day, May 27, and the percentage was 56, which is only slightly above the mean for May at that station. At Madison measurements obtained on five days give a mean of 57 per cent with a maximum of 64 per cent on the 12th. These are only slightly below the corresponding averages for May at Madison.

Table 1.—Solar radiation intensities during May, 1927
[Gram-calories per minute per squre centimeter of normal surface)
Washington, D. C.

: .				St	Sun's zenith distance								
	8 a. m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noon		
Date	75th	Air mass									Local mean		
	time	A. M.					Р. М.				solar time		
	€.	5.0	4.0	3.0	2.0	1 1.0	2.0	3.0	4.0	5.0	e.		
Мау 2	mm. 6.50		cal. 0.74	cal. 0. 92	cal. 1.07		cal.	cal.	cal.	cal.	mm. 5. 36		
May 5	11. 38					1. 27					10. 2		
May 11	13. 61					1. 26							
Маў 20 Маў 27	10. 97 6. 27			7 08	1. 20	1. 34 1. 45	1. 10				9.8		
MRA 41	U. 21	0.00	0.02	U. #0	1. 20	1. 40					1 0. "		
Means Departures		(0. 65) +0. 03	(0. 78) +0. 0 7	(0. 95) +0. 13	(1. 14) +0. 15	1. 30 +0. 02	(1. 10) +0. 11						
	1		,	Madi	son, W	/is.		,		1	1		
May 5	6.02				1, 20						5. 10		
May 12	4.95				1. 26	1.45					4.5		
May 16	4.17				1.20	1.40					3. 1 10. 9		
May 17 May 18	6.02 5.36										6.7		
May 21	12.68										14.6		
May 25	7.04				1.14						7.8		
May 26	8.18					1. 45					6. 70		
				}	1				1		1		
Means] -	1. 20]			
Departures			¦		+0.09	+0.02							
				Linco	ln, Ne	br.							
May 11	4. 57		1.00	1. 13	1, 30 1, 25	1. 54	1. 21	1.02	0. 90		3. 8: 7. 21		
May 13 May 14	8.48 6.02			1.08	1.25	1. 50					4.3		
May 23	13.61					2.30		1.05	0. 92		12.6		
May 25			0.86	1.02	1. 22	1.44					6. 5		
May 28	8.18	ł	0.84			1.44					8.4		
May 30	7. 57					1. 29	1.05	0.85	0.71	0.60	8.8		
Means Departures			0. 90 +0. 09				(1. 13)			(0, 60)			

¹ Extrapolated.

Table 2.—Solar and sky radiation received on a horizontal surface [Gram-calories per square centimeter of norizontal surface]

Week beginning		Ave	Average daily departure from normal						
	Wash- ington	Madi- son	Lin- coln	Chi- cago	New York	Twin Falls	Wash- ington	Madi- son	Lin- coln
1927 Apr. 30 May 7 May 14 May 21 May 28	417 346 375 443 370	473 338 430 366 298	484 415 551 592 438	333 261 261 343 366	304 307 343 296 527	1 616 515 634 (2)	-33 -106 -92 -43 -122	-23 -122 -47 -112 -181	-7 -51 +27 +51 -76
Deficiency	since fir	st of year	on June	3			-7, 602	-6, 048	-6,090

¹ Five-day mean.

POSITIONS AND AREAS OF SUN SPOTS

[Communicated by Capt. Edwin T. Pollock, Superintendent U. S. Naval Observatory]

[Data furnished by Naval Observatory, in cooperation with Harvard, Yerkes, and Mount Wilson observatories]

	Eastern standard civil time		Heliographic		Area t	
Date			Longi- tude	Latitude	Spot	Group
1927 Apr. 1 (Harvard)	л. 13	m. 36	-67. 0 -35. 0	+14.0 +19.0	106	409
Apr. 2 (Harvard)	12	49	+32.0 -73.0 -56.0 -51.0	-23.0 +14.0 +15.0 -23.0	91	91 273 500
Apr. 8 (Harvard)	9	52	-21. 0 -21. 0 +53. 0 -64. 0 -58. 0 -52. 0	+11.0 -12.0 +11.0 +14.0 -16.0 -15.0	182	91 212 121 424
Apr. 9 (Harvard)	9	51	-40.0 +24.0 -51.0 -44.0 -37.0	-13.0 +14.0 +15.0 -15.0 -16.0		288 333 136 106 379
Apr. 13 (Harvard)	10	51	-26.0 +37.0 -2.0 +11.0	-12.0 +14.0 -21.0 -16.0		318 364 76 364
Apr. 25 (Harvard)	9	40	+26.0 -78.0 -31.0 -8.0	-12.0 +4.0 -17.6 -21.0	288 76	61 288
Apr. 29 (Harvard)	16	49	+28.0 -53.0 -19.0	+24.0 -22.0 +4.0	4.5	106 106
Apr. 30 (Harvard)	12	5	+26.0 -16.0 9.0	-14.0 -20.0 +4.0		242 152 136
May 1 (Naval Observatory)	11	48	+36.0 -5.0 +6.5	-12.0 -19.5 +4.5	61	108 108
May 2 (Naval Observatory)	11	45	+50.0 -74.0 +9.5 +19.5	-14.0 +10.0 -19.5 +4.5	123	216 93
May 3 (Naval Observatory)	11	48	+62.5 -60.0 -54.0	-13.5 +10.0 -16.0	77	46 62
May 4 (Harvard)	9	24	+22.5 +33.0 -53.0 -48.0	-19.5 +4.5 -3.0 +9.0	46	247 50
			-41.0 -6.0 +34.0 +43.0	-16.0 +22.0 -19.0 +6.0	49	234 484 63
May 5 (Naval Observatory)	11	44	-38, 5 -26, 0 +26, 0 +44, 5	-8.0 -16.5 +22.0 -23.0	19	139 201 62
May 6 (Harvard)	9	2	+53.0 -28.0 -23.0 -15.0	-18.0 -6.0 -10.0 -16.0	83	247 95 885
May 7 (Naval Observatory)	12	25	+35.0 +86.0 -10.0 +3.0 +14.0	+24.0 +29.0 -7.5 -14.5 +19.0	68	118 108 216 77
1 Areas are corrected for foreshortenin	o and	974 (+50.5 +84.0	+23. 5 -18. 0	15	247

Areas are corrected for foreshortening and are expressed in millionths of sun's visible hemisphere.

¹ Main-spring of register clock broken.